

## GH とストレス・麻酔

KW (10/02/13)

5 分間の寒冷暴露で有意に低下した (43%に). (1)

5 分間の緊縛ストレス (immobilization) で有意に低下した. (1)

毎日 8 時間, 10 日間の immobilization で低下した. (2)

この状態で GnRH+TRH の投与で更に低下した.

絶食で増加した. (3)

Ketamine(100mg/kg)+Xylazine(10mg/kg)のと紆余で GH レベルが上昇した. (4)

### 参考文献

1. Specific hormonal and neurochemical response to different stresses.  
Lenox RH, Kant GJ, Sissions GR., et al.  
Neuroendocrinology 30:300-308, 1980
2. Effects of chronic immobilization stress on pituitary hormone secretion, on hypothalamic factor levels, and on pituitary responsiveness to LHRH and TRH in female rats.  
Du Ruisseau P, Tache Y, Brazeau P, and Collu R.  
Neuroendocrinology 29: 90-99, 1979
3. Changes in serum levels of gonadotropins and testosterone in the male rat in response to fasting, surgery and ether.  
Howland BE, Beaton DB and Jack MI.  
Experientia 30:1223-1225, 1974
4. Acute hyperglycemia induced by ketamine/xylazine anesthesia in rats: Mechanism and implication for preclinical models.  
Saha JK, Xia J, Grondin JM, et al.  
Exp Biol Med (Maywood) 230: 777-784, 2005